

Numerical exploration
in sphere packing,
Fourier analysis, and physics


Lecture 4

Henry Cohn

PCM1 2022

Does E_8 minimize energy for every convex, decreasing potential f ?

No:



$$\text{Energy of } E_8: 240 f(\sqrt{2}) + 2160 f(\sqrt{4}) + \dots \\ = 240$$

$$\text{Energy of } \mathbb{Z}^8: 16 f(1) + 112 f(\sqrt{2}) + 448 f(\sqrt{3}) \\ + \dots \\ = 152$$

$$152 < 240$$

Today:


We'll use modular forms to interpolate a function.

When I first learned about modular forms, they were really intimidating.

If you're not already comfortable with them, don't worry. Lots of things are surprisingly concrete.

← fundamental domain for $SL_2(\mathbb{Z})$

Modular Forms



This is not our fate!

$SL_2(\mathbb{R})$ acts on the upper
half-plane

$$\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$$

by linear fractional
transformations,

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R})$$

$$z \in \mathbb{H}$$

$$\gamma z = \frac{az + b}{cz + d} \in \mathbb{H}$$


This is the conformal
automorphism group of \mathbb{H} .

(also orientation-preserving
isometries of the hyperbolic plane)

$SL_2(\mathbb{Z})$ is an important discrete subgroup of $SL_2(\mathbb{R})$.

\mathbb{H}^1 's generated by
 $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad z \mapsto -1/z$

and $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad z \mapsto z+1$



A modular form for $SL_2(\mathbb{Z})$ of weight k is a holomorphic function $f: \mathbb{H} \rightarrow \mathbb{C}$

such that

$$(1) \quad f|_k \gamma = f \quad \text{for all}$$

where $\gamma \in SL_2(\mathbb{Z})$,

$$(f|_k \gamma)(z) = (cz+d)^{-k} f\left(\frac{az+b}{cz+d}\right)$$

$$\text{if } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

and (2) $f(z)$ remains bounded as $\text{Im } z \rightarrow \infty$
("holomorphic at the cusp")

Why?

This theory is justified by examples and applications. It's not obviously compelling to a beginner.

These objects can be understood at many levels. For example, $f(z) dz^{\otimes k/2}$ is a section of the $k/2$ tensor power of the cotangent bundle of $SL_2(\mathbb{Z}) \backslash \mathbb{H}$.

(But this just raises the question of why we want such sections.)

In terms of S and T ,

$$f(-1/z) = z^k f(z)$$

$$f(z+1) = f(z).$$

These functional equations
come up surprisingly often.

The former is often deeper.

Example: theta series

$$\theta_{\Lambda}(z) = \sum_{x \in \Lambda} e^{\pi i |x|^2 z}$$

converges for $z \in \mathbb{H}$

Think of it as a generating function for the vector lengths.

For $\Lambda = E_8$, $|x|^2 \in 2\mathbb{Z}$ for all x .

$$\theta_{E_8}(z) = \sum_{n=0}^{\infty} N_n e^{2\pi i n z}$$

where $N_n = \#\{x \in E_8 : |x|^2 = 2n\}$

$$\begin{aligned} \Theta_{E_8}(z) &= \sum_{x \in E_8} e^{\pi i |x|^2 z} \\ &= \sum_{n=0}^{\infty} N_n e^{z \pi i n z} \end{aligned}$$

This implies $\Theta_{E_8}(z+1) = \Theta_{E_8}(z)$.

For S , we use Poisson summation.

$$E_8^* = E_8$$

Fourier transform
of
 $x \mapsto e^{\pi i z |x|^2}$
 $y \mapsto \left(\frac{i}{z}\right)^{d/2} e^{\pi i (-1/z) |x|^2}$
 in \mathbb{R}^d
 Here $d=8$, $i^{d/2} = 1$.
 Looks like S !

Poisson:

$$\sum_{x \in E_8} e^{\pi i z |x|^2} = z^{-4} \sum_{y \in E_8} e^{\pi i (-y) |y|^2}$$

$$\Theta_{E_8}(z) = z^{-4} \Theta_{E_8}(-y(z))$$

So

$$\Theta_{E_8} |_4 S = \Theta_{E_8}$$

$$\Theta_{E_8} |_4 T = \Theta_{E_8}$$

$$\Theta_{E_8}(z) = 1 + 240e^{2\pi i z} + \dots$$

$\rightarrow 1$ as $\text{Im } z \rightarrow \infty$

modular form of weight 4

So our motivation is that
we want to understand
functions like the theta
series of E_8 (weight 4)
or Λ_{24} (weight 12).

Eisenstein series

(unfortunate notational clash w/ E_g)

$$E_k(z) = \frac{1}{2\zeta(k)} \sum_{\substack{(m,n) \in \mathbb{Z}^2 \\ (m,n) \neq (0,0)}} \frac{1}{(mz+n)^k}$$

(just a
normalizing
factor)

$$\begin{aligned} (m,n) &\in \mathbb{Z}^2 \\ (m,n) &\neq (0,0) \end{aligned}$$

converges absolutely for $k \geq 2$
vanishes when k is odd

$$E_k(z+i) = E_k(z)$$

$$E_k(-1/z) = z^k E_k(z)$$

(just from manipulating
the series)

modular form of weight k
(for $k \geq 2$)

Key fact: E_4 and E_6 generate
the graded ring of modular
forms for $SL_2(\mathbb{Z})$

$$\text{E.g., } \Theta_{E_8} = E_4$$

$$\Theta_{\Lambda_{24}} = \frac{7}{12} E_4^3 + \frac{5}{12} E_6^2$$

There are many other beautiful
modular forms, e.g., Ramanujan's
discriminant

$$\Delta = \frac{E_4^3 - E_6^2}{1728}$$

Thanks to invariance under $z \mapsto z+1$, these functions all have Fourier series.

$$E_4(z) = 1 + 240 \sum_{n \geq 1} \sigma_3(n) e^{2\pi i n z}$$

$$E_6(z) = 1 - 504 \sum_{n \geq 1} \sigma_5(n) e^{2\pi i n z}$$

where $\sigma_k(n) = \sum_{d|n} d^k$.



It's conventional to set $q = e^{2\pi i z}$.

$$\Delta(z) = q \prod_{n \geq 1} (1 - q^n)^{24}$$

vanishes at ∞ , but

$$\Delta(z) \neq 0 \text{ for } z \in \mathbb{H}.$$

Plot of $E_4(z)$ for $-1 \leq \operatorname{Re} z \leq 1$
 $0 < \operatorname{Im} z \leq 1$
(following Lowry - Duda)



(decorated w/ hyperbolic tiling)

We'll also need modular forms for the subgroup

$$\Gamma(2) = \left\{ \gamma \in SL_2(\mathbb{Z}) : \gamma \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{2} \right\}.$$

(i.e., $f|_k \gamma = f$ for $\gamma \in \Gamma(2)$, and $(f|_k \gamma)(z)$ bounded as $\text{Im} z \rightarrow \infty$ for all $\gamma \in SL_2(\mathbb{Z})$)

Generated by weight 2 forms U, W :

$$U = (\Theta_{\mathbb{Z}})^4$$

$$W = U|_2 T$$

$$V = U - W$$

(many notations used)

$$U|_2 T = W$$

$$V|_2 T = -V$$

$$W|_2 T = U$$

$$U|_2 S = -U$$

$$V|_2 S = -W$$

$$W|_2 S = -V$$

Where are we going with this?

Let's interpolate radial Schwartz functions on \mathbb{R}^8 .

We'll do 1st order interpolation

for simplicity: recover f from

$f(\sqrt{n})$ and $\hat{f}(\sqrt{n})$ w/ $n \geq 2$.

(Radchenko and Viazovska, 2019)

We'll build only one function in

an interpolation basis, as a

taste.

We'll also look at -1 eigenfunctions

of the Fourier transform.

Fourier eigenfunctions:

$$\hat{\hat{f}}(x) = f(-x) \text{ by}$$

Fourier inversion,

$$\text{so } \hat{\hat{f}} = f \text{ for radial}$$

(or any even) f .

We can write

$$f = \underbrace{\frac{f - \hat{f}}{2}}_{-1 \text{ eigenfunction}} + \underbrace{\frac{f + \hat{f}}{2}}_{+1 \text{ eigenfunction}}$$

These cases turn out to require somewhat different techniques.

We'll focus on -1 eigenfunctions.

Goal:

Construct a radial Schwartz
function $g: \mathbb{R}^8 \rightarrow \mathbb{R}$

such that

$$\hat{g} = -g$$


$$g(\sqrt{2}) = 1$$

$$g(\sqrt{n}) = 0 \quad \text{for } n \geq 3$$

(If we rotate the position of the
1 and analyze both eigenvalues
 ± 1 , we'll get an interpolation
basis.)

Construction:

$$g(x) = \frac{1}{2} \int_{-1}^1 \psi(z) e^{\pi i z |x|^2} dz$$



Which properties for ψ
will make this work?

$$g(x) = \frac{1}{2} \int_{-1}^1 \psi(z) e^{\pi i z |x|^2} dz$$

Let's suppose $\psi(z+2) = \psi(z)$.

Fourier series

$$\psi(z) = \sum_{n \in \mathbb{Z}} a_n e^{\pi i n z}$$

$$g(\sqrt{n}) = \frac{1}{2} \int_{-1}^1 \psi(z) e^{\pi i n z} dz$$

$$= a_{-n} \text{ by orthogonality}$$

We want $a_{-2} = 1$, $a_n = 0$ for
 $n < -2$.

$$g(x) = \frac{1}{2} \int_{-1}^1 \psi(z) e^{\pi i z |x|^2} dz$$

$$\psi(z) = e^{-2\pi i z} + \sum_{n \geq -1} a_n e^{\pi i n z}$$

$\psi(z)$ should vanish as $z \rightarrow \pm 1$ to get a Schwartz function

When is g an eigenfunction of the Fourier transform?

$$g(x) = \frac{1}{2} \int_{-1}^1 \psi(z) e^{\pi i z |x|^2} dz$$

8-dim'l Fourier transform of
 $x \mapsto e^{\pi i z |x|^2}$

is $x \mapsto z^{-4} e^{\pi i (-1/z) |x|^2}$

so

$$\hat{g}(x) = \frac{1}{2} \int_{-1}^1 \psi(z) z^{-4} e^{\pi i (-1/z) |x|^2} dz$$

$$\hat{g}(x) = \frac{1}{2} \int_{-1}^1 \psi(z) z^{-4} e^{\pi i (-1/2) |x|^2} dz$$

let $u = -1/z$.

$$= \frac{1}{2} \int_1^{-1} \psi(-1/u) u^4 e^{\pi i u |x|^2} \frac{du}{u^2}$$

$$= -\frac{1}{2} \int_{-1}^1 \underbrace{u^2 \psi(-1/u)}_{(\psi|_S)(u)} e^{\pi i u |x|^2} du$$

so $\hat{g} = -g$ if

$$\psi|_S = \psi$$

So ψ should satisfy:

- $\psi|_{-2} T^2 = \psi$ (i.e., $\psi(z+2) = \psi(z)$)
- $\psi|_{-2} S = \psi$

• $\psi(z) = e^{-2\pi i z} + \sum_{n \geq -1} a_n e^{\pi i n z}$

- ψ vanishes at ± 1

Note: only meromorphic at ∞
and negative weight,

but $\Delta \cdot \psi$ has weight 10
and is holomorphic at ∞ .

$\Delta \cdot \psi$ is modular for $\Gamma(2)$
(a subgroup of the group
generated by S, T^2)

linear combination of
 $U^5, U^4W, U^3W^2, U^2W^3, UW^4, W^5$

and we know how S, T act on
 U, W

Simultaneous linear equations
w/ unique sol'n

$$\psi = \frac{2U^4W - U^5}{\Delta}$$

$$\psi = \frac{2U^4 w - U^5}{\Delta}$$

$$= q^{-1} + 8q^{-1/2} - 240 - 6176q^{1/2} - \dots$$

$$\implies g(\sqrt{n}) = \begin{cases} -240 & n=0 \\ 8 & n=1 \\ 1 & n=2 \\ 0 & n \geq 3 \end{cases}$$

as desired

What's left?

- a somewhat more elaborate integral for 2nd order interpolation
- using modular forms for $\Gamma(2)$, quasimodular form E_2 , and $\log \lambda$

This suffices to get each interpolation basis function.

Then

- systematize to solve for generating functions
- prove completeness