








Observation of the Yamaji effect in a cuprate and FL^* theory of the pseudogap

Simons Ultra-Quantum Matter Collaboration Meeting
Institute for Advanced Study, Princeton
October 20, 2025

Subir Sachdev









1-p electrons

1+p holes

Luttinger, 1960: Area enclosed by the Fermi surface is the same as that for free fermions *with the same symmetry*.





$1-p$ electrons

$1+p$ holes

Luttinger, 1960: Area enclosed by the Fermi surface is the same as that for free fermions *with the same symmetry*.
Oshikawa, 2000: Area constrained by a 't Hooft anomaly of global U(1) and translations

Keimer, Kivelson, Norman, Uchida, and Zaanen, *Nature* **518**, 179 (2015)

Kyle M. Shen *et al.*, *Science* **307**, 901 (2005)



‘Fermi arcs’ ?


T. Senthil, S. S., M.Vojta, PRL **90**, 216403 (2003)

Observation of the Yamaji effect in a cuprate superconductor


Mun K. Chan¹✉, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

nature physics
arXiv:2411.10631


Published online: 16 September 2025




$\text{HgBa}_2\text{CuO}_{4+\delta}$
(Hg1201)



$\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$
(YBCO)



$\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$
(LSCO)






Observation of the Yamaji effect in a cuprate superconductor

nature physics

arXiv:2411.10631

Mun K. Chan¹, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

Published online: 16 September 2025






Observation of the Yamaji effect in a cuprate superconductor

nature physics

arXiv:2411.10631

Mun K. Chan¹✉, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

Published online: 16 September 2025



At the Yamaji angle, the orbits in the plane orthogonal to $\mathbf{B}$ have an area which is independent of momentum in the c direction, to first order in the hopping along the c direction.

K.Yamaji JPSJ **58**, 1520 (1989)




Observation of the Yamaji effect in a cuprate superconductor

nature physics

arXiv:2411.10631

Mun K. Chan¹, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

Published online: 16 September 2025




Doping
 $p = 0.1$

“The small size of the pockets determined from the Yamaji effect is ... approximately 1.3% of the Brillouin zone area”


***Metals obtained by
doping Mott insulators***

Ordinary metal




At large p , we obtain a gas of nearly free fermionic holes of density $1+p$ (relative to the filled band with 2 electrons per site)

Luttinger area.
No broken symmetry




Area $(1+p)/2$

Insulating antiferromagnet




Doping an insulating antiferromagnet with holes of density p

AF metal



Luttinger area.
Broken symmetry





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Spin liquid with density p of spinless, charge $+e$ 'holons'. No coherent inter-layer transport, so no Yamaji effect.



Area $p/4$



T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003);
R. K. Kaul, A. Kolezhuk, M. Levin, S. S., T. Senthil, PRB **75**, 235122 (2007);
R. K. Kaul, Y. B. Kim, S. S., T. Senthil, Nature Physics **4**, 28 (2008)

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Spin liquid with density p of spinless, charge $+e$ 'holons'. No coherent inter-layer transport, so no Yamaji effect.



Area $p/4$



T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003);
R. K. Kaul, A. Kolezhuk, M. Levin, S. S., T. Senthil, PRB **75**, 235122 (2007);
R. K. Kaul, Y. B. Kim, S. S., T. Senthil, Nature Physics **4**, 28 (2008)

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Spin liquid with density p of spinless, charge $+e$ 'holons'. No coherent inter-layer transport, so no Yamaji effect.



Area $p/4$



T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003);
R. K. Kaul, A. Kolezhuk, M. Levin, S. S., T. Senthil, PRB **75**, 235122 (2007);
R. K. Kaul, Y. B. Kim, S. S., T. Senthil, Nature Physics **4**, 28 (2008)

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Spin liquid with density p of spinless, charge $+e$ 'holons'. No coherent inter-layer transport, so no Yamaji effect.



Area $p/4$



T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003);
R. K. Kaul, A. Kolezhuk, M. Levin, S. S., T. Senthil, PRB **75**, 235122 (2007);
R. K. Kaul, Y. B. Kim, S. S., T. Senthil, Nature Physics **4**, 28 (2008)

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Spin liquid with density p of spinless, charge $+e$ 'holons'. No coherent inter-layer transport, so no Yamaji effect.



Area $p/4$



T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003);
R. K. Kaul, A. Kolezhuk, M. Levin, S. S., T. Senthil, PRB **75**, 235122 (2007);
R. K. Kaul, Y. B. Kim, S. S., T. Senthil, Nature Physics **4**, 28 (2008)

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied
by sum of spin liquid (1) and
Fermi surface anomalies (p)

Also has
charge 0,
spin-1/2
'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Also has charge 0, spin-1/2 'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Also has charge 0, spin-1/2 'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Also has charge 0, spin-1/2 'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Also has charge 0, spin-1/2 'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied
by sum of spin liquid (1) and
Fermi surface anomalies (p)

Also has
charge 0,
spin-1/2
'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied
by sum of spin liquid (1) and
Fermi surface anomalies (p)

Also has
charge 0,
spin-1/2
'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

Holon metal

Oshikawa anomaly is satisfied
by sum of spin liquid (1) and
Fermi surface anomalies (p)

Also has
charge 0,
spin-1/2
'spinons'





Area $p/4$

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied
by sum of spin liquid (1) and
Fermi surface anomalies (p)

Metal with
density p of
spin-1/2,
charge $+e$
'holes' (or
'magnetic
polarons')
and
charge 0,
spin-1/2
'spinons'



Area $p/8$

$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \begin{array}{|c|} \hline \bullet \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \begin{array}{|c|} \hline \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \begin{array}{|c|} \hline \bullet \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

$$\text{Blue oval} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \text{Green rectangle} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \begin{array}{|c|} \hline \bullet \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \begin{array}{|c|} \hline \bullet \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



Area $p/8$

$$\text{Blue oval} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \text{Green rectangle} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \begin{array}{|c|} \hline \bullet \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



$$\begin{array}{|c|c|} \hline \bullet \quad \bullet & = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \\ \hline \end{array} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)



M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'





$$\text{blue oval} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \text{green rectangle} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)


M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

Doping an insulating antiferromagnet with holes of density p




Area $p/4$

AF metal and SDW fluctuation




$$\text{blue ellipse} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2}$$

Holon metal




Area $p/4$



FL*

$$\text{green rectangle} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$




Area $p/8$

Quantization of spin liquid anomaly implies Fermi surface areas are also quantized and robust to all corrections.

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003);
 R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)
 M. Punk, A. Allais, and S. S., PNAS **112**, 9552 (2015)
 E. Mascot, A. Nikolaenko, M. Tikhanovskaya, Ya-Hui Zhang, D. K. Morr, S. S., PRB **105**, 075146 (2022)

Hole dynamics in an antiferromagnet across a deconfined quantum critical point

Ribhu K. Kaul,¹ Alexei Kolezhuk,^{1,2} Michael Levin,¹ Subir Sachdev,¹ and T. Senthil^{3,4}



The dashed line in the Néel phase indicates the boundary of the magnetic Brillouin zone. Only the Fermi surfaces within this zone contribute to the Luttinger counting, and so the area of each ellipse is $\mathcal{A}_F = (2\pi)^2 \delta/4$. In the VBS phase, all four pockets are inequivalent, and so the area of each ellipse is $\mathcal{A}_F = (2\pi)^2 \delta/8$.

Factor of 2 between
SDW fluctuation
and FL*




Observation of the Yamaji effect in a cuprate superconductor

nature physics

arXiv:2411.10631

Mun K. Chan¹, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

Published online: 16 September 2025



Doping
 $p = 0.1$

“The small size of the pockets determined from the Yamaji effect is ... approximately 1.3% of the Brillouin zone area”




Observation of the Yamaji effect in a cuprate superconductor

nature physics

arXiv:2411.10631

Mun K. Chan¹, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

Published online: 16 September 2025



Doping
 $p = 0.1$

“The small size of the pockets determined from the Yamaji effect is ... approximately 1.3% of the Brillouin zone area”

FL* pocket fraction = $p/8 = 1.25\%$!

Fluctuating AF metal fraction = $p/4 = 2.5\%$.

Jing-Yu Zhao, S. Chatterjee, S. S.,
and Ya-Hui Zhang, arXiv:2510.13943

($p/8$ also in YRZ ansatz, Peter Johnson photoemission,
and Jenny Hoffman and Seamus Davis STMs; Stanescu-Kotliar)



Observation of the Yamaji effect in a cuprate superconductor

nature physics

arXiv:2411.10631

Mun K. Chan¹, Katherine A. Schreiber¹, Oscar E. Ayala-Valenzuela¹,
Eric D. Bauer², Arkady Shekhter¹ & Neil Harrison¹

Published online: 16 September 2025



Doping
 $p = 0.1$

“The small size of the pockets determined from the Yamaji effect is ... approximately 1.3% of the Brillouin zone area”

FL* pocket fraction = $p/8 = 1.25\%$!

Fluctuating AF metal fraction = $p/4 = 2.5\%$.

Jing-Yu Zhao, S. Chatterjee, S. S.,
and Ya-Hui Zhang, arXiv:2510.13943


($p/8$ also in Peter Johnson photoemission,
and Jenny Hoffman and Seamus Davis STMs; Stanescu-Kotliar)

Coupling hole pockets to spin liquid

(the spin liquid is required
to satisfy Oshikawa anomaly)



Maine Christos
Caltech



Pietro Bonetti



Alexander
Nikolaenko

Thermal $SU(2)$ lattice gauge theory of
the cuprate pseudogap: reconciling
Fermi arcs and hole pockets

H. Pandey, M. Christos, P.M. Bonetti,
R. Shanker, A. Nikolaenko, S. Sharma,
S.S., arXiv:2507.05336

The Institute of
Mathematical
Sciences,
Chennai



Harshit Pandey




Ravi Shanker



Sayantan Sharma



Maine Christos
Caltech



Pietro Bonetti



Alexander
Nikolaenko



Aavishkar Patel
ICTS, Bengaluru

arXiv > cond-mat > arXiv:2508.20164

Condensed Matter > Strongly Correlated Electrons

[Submitted on 27 Aug 2025]



Critical quantum liquids and the cuprate high temperature superconductors

Doping an insulating antiferromagnet with holes of density p

FL*

Oshikawa anomaly is satisfied by sum of spin liquid (1) and Fermi surface anomalies (p)

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'



$$\text{blue oval} = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle) / \sqrt{2} \quad \text{green rectangle} = (|\uparrow\circ\rangle + |\circ\uparrow\rangle) / \sqrt{2}$$

Area $p/8$

T. Senthil, S. S., M. Vojta, PRL **90**, 216403 (2003); R. K. Kaul, A. Kolezhuk, M. Levin, S.S., T. Senthil, PRB **75**, 235122 (2007)

M. Punk, A. Allais, and S. Sachdev, PNAS **112**, 9552 (2015)

1. Dope spin liquid with *holes*, not holons.

The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.


Ya-Hui Zhang and S. S., PRR **2**, 023172 (2020)

1. Dope spin liquid with *holes*, not holons.

The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.

2. Hole pockets are obtained by hybridizing the large Fermi surface of $1 - p$ electrons c with the Fermi surface of 1 ancilla fermion f_1 . The hybridization Φ also determines the electronic gap near $(\pi, 0)$ (similar to electrons in Weinberg-Salam theory).

$$H_{\text{hole pockets}} = \sum_{i,j} \left[-t_{ij} c_{i\alpha}^\dagger c_{j\alpha} - t_{1,ij} f_{1i\alpha}^\dagger f_{1j\alpha} \right] - \sum_i \Phi (c_{i\alpha}^\dagger f_{1i\alpha} + f_{1i\alpha}^\dagger c_{i\alpha})$$



Ya-Hui Zhang and S. S., PRR **2**, 023172 (2020)


Kai-Yu Yang, T. M. Rice, Fu-Chun Zhang, PRB **73**, 174501 (2006)

Y. Qi and S. S., PRB **81**, 115129 (2010)

1. Dope spin liquid with *holes*, not holons.
The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.
2. Hole pockets are obtained by hybridizing the large Fermi surface of $1 - p$ electrons c with the Fermi surface of 1 ancilla fermion f_1 . The hybridization Φ also determines the electronic gap near $(\pi, 0)$
(similar to electrons in Weinberg-Salam theory).
3. Building on recent numerical evidence, we choose the π -flux spin liquid with fermionic spinons f coupled to an emergent SU(2) gauge field U
(similar to neutrinos in Weinberg-Salam theory).

1. Dope spin liquid with *holes*, not holons.
The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.
2. Hole pockets are obtained by hybridizing the large Fermi surface of $1 - p$ electrons c with the Fermi surface of 1 ancilla fermion f_1 . The hybridization Φ also determines the electronic gap near $(\pi, 0)$
(similar to electrons in Weinberg-Salam theory).
3. Building on recent numerical evidence, we choose the π -flux spin liquid with fermionic spinons f coupled to an emergent SU(2) gauge field U
(similar to neutrinos in Weinberg-Salam theory).


$$H_{\text{spin liquid}} = iJ \sum_{\langle ij \rangle} e_{ij} \left(\Psi_i^\dagger U_{ij} \Psi_j - \Psi_j^\dagger U_{ji} \Psi_i \right); \quad \Psi_i = \begin{pmatrix} f_{i\uparrow} \\ f_{i\downarrow} \end{pmatrix}$$




1. Dope spin liquid with *holes*, not holons.
The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.
2. Hole pockets are obtained by hybridizing the large Fermi surface of $1 - p$ electrons c with the Fermi surface of 1 ancilla fermion f_1 . The hybridization Φ also determines the electronic gap near $(\pi, 0)$
(similar to electrons in Weinberg-Salam theory).
3. Building on recent numerical evidence, we choose the π -flux spin liquid with fermionic spinons f coupled to an emergent SU(2) gauge field U
(similar to neutrinos in Weinberg-Salam theory).

Low energy: SU(2) gauge theory with $N_f = 2$
massless Dirac spinons in 2+1 D.

(QCD is a SU(3) gauge theory with $N_f = 3$ massless Dirac quarks in 3+1 D.)



1. Dope spin liquid with *holes*, not holons.
The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.
2. Hole pockets are obtained by hybridizing the large Fermi surface of $1 - p$ electrons c with the Fermi surface of 1 ancilla fermion f_1 . The hybridization Φ also determines the electronic gap near $(\pi, 0)$
(similar to electrons in Weinberg-Salam theory).
3. Building on recent numerical evidence, we choose the π -flux spin liquid with fermionic spinons f coupled to an emergent SU(2) gauge field U
(similar to neutrinos in Weinberg-Salam theory).
Low energy: SU(2) gauge theory with $N_f = 2$
massless Dirac spinons in 2+1 D.
(QCD is a SU(3) gauge theory with $N_f = 3$ massless Dirac quarks in 3+1 D.)



1. Dope spin liquid with *holes*, not holons.
The Ancilla Layer Model (ALM) enables a theory of FL* with hole pockets and a general spin liquid.
2. Hole pockets are obtained by hybridizing the large Fermi surface of $1 - p$ electrons c with the Fermi surface of 1 ancilla fermion f_1 . The hybridization Φ also determines the electronic gap near $(\pi, 0)$
(similar to electrons in Weinberg-Salam theory).
3. Building on recent numerical evidence, we choose the π -flux spin liquid with fermionic spinons f coupled to an emergent SU(2) gauge field U
(similar to neutrinos in Weinberg-Salam theory).
4. Couple hole pockets coupled to critical spin liquid by a charge e , SU(2) fundamental Higgs field B
(similar to Yukawa coupling of Higgs field in Weinberg-Salam theory).

- To create the electron operator from the spinon Ψ , we have to fuse it with a charge e , gauge SU(2) fundamental, spinless boson $B_i = (B_{1i}, B_{2,i})$, whose symmetries follow from those of Ψ .

This yields the quadratic and quartic energy functionals $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$

$$\mathcal{E}_2[B, U] = r \sum_i B_i^\dagger B_i + iw \sum_{\langle ij \rangle} e_{ij} \left(B_i^\dagger U_{ij} B_j - B_j^\dagger U_{ji} B_i \right)$$


$$+ \kappa \sum_{\square} \left\{ 1 - \frac{1}{2} \text{ReTr} \prod_{ij \in \square} U_{ij} \right\}$$

$$H_{\text{Yukawa}} = ig \sum_i \left(B_{1i}^* f_{1i\sigma}^\dagger f_{i\sigma} + B_{2i}^* \varepsilon_{\sigma\sigma'} f_{1i\sigma}^\dagger f_{i\sigma'}^\dagger - \text{H.c.} \right)$$

$$\begin{aligned}
H_{\text{hole pockets}} &= \sum_{i,j} \left[-t_{ij} c_{i\alpha}^\dagger c_{j\alpha} - t_{1,ij} f_{1i\alpha}^\dagger f_{1j\alpha} \right] \\
&\quad - \sum_i \Phi \left(c_{i\alpha}^\dagger f_{1i\alpha} + f_{1i\alpha}^\dagger c_{i\alpha} \right) \\
H_{\text{spin liquid}} &= iJ \sum_{\langle ij \rangle} e_{ij} \left(\Psi_i^\dagger U_{ij} \Psi_j - \Psi_j^\dagger U_{ji} \Psi_i \right); \quad \Psi_i = \begin{pmatrix} f_{i\uparrow}^\dagger \\ f_{i\downarrow}^\dagger \end{pmatrix} \\
\mathcal{E}[B, U] &= r \sum_i B_i^\dagger B_i + iw \sum_{\langle ij \rangle} e_{ij} \left(B_i^\dagger U_{ij} B_j - B_j^\dagger U_{ji} B_i \right) \\
&\quad + \mathcal{E}_4[B, U] + \kappa \sum_{\square} \left\{ 1 - \frac{1}{2} \text{ReTr} \prod_{ij \in \square} U_{ij} \right\} \\
H_{\text{Yukawa}} &= ig \sum_i \left(B_{1i}^* f_{1i\sigma}^\dagger f_{i\sigma} + B_{2i}^* \varepsilon_{\sigma\sigma'} f_{1i\sigma}^\dagger f_{i\sigma'}^\dagger - \text{H.c.} \right)
\end{aligned}$$


} Hole pockets
 } Spinons
 } Higgs potential
 } Yukawa coupling

The pseudogap metal
and its transition to
d-wave superconductivity





Hole pockets and spin liquid.
 If the spin liquid is stable at $T = 0$,
 hole pockets remain intact
 despite Yukawa coupling to spin liquid.

- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T > T_c$.
Diagonalize Hamiltonian for electrons and spinons for each realization of B, U .





- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T > T_c$.
Diagonalize Hamiltonian for electrons and spinons for each realization of B, U .



Kyle M. Shen, ... Z.-X. Shen, Science **307**, 901 (2005)

H. Pandey, M. Christos, P.M. Bonetti, R. Shanker, A. Nikolaenko, S. Sharma, S.S., arXiv:2507.05336


- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T > T_c$.
Diagonalize Hamiltonian for electrons and spinons for each realization of B, U .



Fermi arcs
co-exist with
quantum
oscillations
of hole pockets
of area $p/8$.

- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T \rightarrow 0$.

V_{Higgs} chosen so that the ground state is a d -wave superconductor,
and second best state is a period-4 stripe.




$T = 0$ phase diagram

H. Pandey,
M. Christos,
P.M. Bonetti,
R. Shanker,
A. Nikolaenko,
S. Sharma,
S.S.,
arXiv:2507.05336

Minimize V_{Higgs} w.r.t. $\left\langle \begin{pmatrix} B_{1i} \\ B_{2i} \end{pmatrix} \right\rangle$
Set $U_{ij} = 1$.


- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T \rightarrow 0$.




H. Pandey,
M. Christos,
P.M. Bonetti,
R. Shanker,
A. Nikolaenko,
S. Sharma,
S.S.,
arXiv:2507.05336

See also
Jia-Xin Zhang
and S. S.,
PRB **110**,
235120
(2024)

- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T \rightarrow 0$.




Bond density



Phase of pairing amplitude

H. Pandey,
M. Christos,
P.M. Bonetti,
R. Shanker,
A. Nikolaenko,
S. Sharma,
S.S.,
arXiv:2507.05336

See also
Jia-Xin Zhang
and S. S.,
PRB **110**,
235120
(2024)




A Four Unit Cell Periodic Pattern of Quasi-Particle States Surrounding Vortex Cores in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$

J. E. Hoffman, E. W. Hudson,
K. M. Lang, V. Madhavan,
H. Eisaki, S. Uchida, J.C. Davis
Science **295**, 466 (2002)

- Born-Oppenheimer theory of thermal fluctuations:
Classical thermal ensemble of $\mathcal{E}_2[B, U] + \mathcal{E}_4[B, U]$ at a temperature $T \lesssim T_{KT}$.

$\Upsilon =$
Helicity
Modulus




H. Pandey,
M. Christos,
P.M. Bonetti,
R. Shanker,
A. Nikolaenko,
S. Sharma,
S.S.,
arXiv:2507.05336

Doping an insulating antiferromagnet with holes of density p

FL*

Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'


The d -wave superconductor is obtained by confining the SU(2) gauge field, and this exploits the pre-existing pairing in the blue dimers. Such a state is smoothly connected to the BCS d -wave superconductor.



Doping an insulating antiferromagnet with holes of density p

FL*


Metal with density p of spin-1/2, charge $+e$ 'holes' (or 'magnetic polarons') and charge 0, spin-1/2 'spinons'




The d -wave superconductor is obtained by confining the SU(2) gauge field, and this exploits the pre-existing pairing in the blue dimers. Such a state is smoothly connected to the BCS d -wave superconductor.

Applying BCS pairing to the hole pockets (green dimers) yields a distinct SC* state, which is *not* smoothly connected to the BCS d -wave state.

The cuprate phase diagram




Quantum-criticality
of a quantum phase transition
between two metals
(FL^* and FL) at $p = p_c$,
with no symmetry breaking.
 Φ was the hybridization
between c and f_1 in theory of
hole pockets—it is
now a Higgs field.



Explain with a local, two-dimensional extension of the Sachdev-Ye-Kitaev (SYK) model of mobile electrons, the 2D-Yukawa-SYK model with a spatially random Yukawa coupling between the large Fermi surface and Φ : a critical charge liquid


Aavishkar A. Patel, Haoyu Guo, Ilya Esterlis, S. S., *Science* **381**, 790 (2023)
 Chenyuan Li, Aavishkar A. Patel, Haoyu Guo, Davide Valentini, Jorg Schmalian, S.S., Ilya Esterlis, *PRL* **133**, 186502 (2024)



Quantum-criticality
of a quantum phase transition
between two metals
(FL* and FL) at $p = p_c$,
with no symmetry breaking.
 Φ was the hybridization
between c and f_1 in theory of
hole pockets—it is
now a Higgs field.


Both metals lead to the same
 d -wave superconductor at lower
temperatures, and so there is
transition at $p = p_c$ within the
superconducting state.

M. Christos, Zhu-Xi Luo, L. Shackleton, Ya-Hui Zhang,
M. S. Scheurer, and S. S., PNAS **120**, e2302701120 (2023)



Quantum-criticality
of a quantum phase transition
between two metals
(FL* and FL) at $p = p_c$,
with no symmetry breaking.
 Φ was the hybridization
between c and f_1 in theory of
hole pockets—it is
now a Higgs field.

Both metals lead to the same
 d -wave superconductor at lower
temperatures, and so there is
transition at $p = p_c$ within the
superconducting state.



Quantum-criticality
of a quantum phase transition
between two metals
(FL* and FL) at $p = p_c$,
with no symmetry breaking.
 Φ was the hybridization
between c and f_1 in theory of
hole pockets—it is
now a Higgs field.

Both metals lead to the same
 d -wave superconductor at lower
temperatures, and so there is
transition at $p = p_c$ within the
superconducting state.