



Rigidity Phenomena in random point sets and Applications

Subhro Ghosh
Princeton University




Poisson
Process

Background




Poisson
Process




Ginibre
Ensemble


Background



Poisson
Process



Ginibre
Ensemble



Gaussian
Zeroes

- Finite n : $\mu_n =$ Eigenvalues of $G_n = ((\xi_{ij}))_{1 \leq i, j \leq n}$, ξ_{ij} i.i.d $N_{\mathbb{C}}(0, 1)$ (NO normalization by \sqrt{n})

- Finite n : $\mu_n =$ Eigenvalues of $G_n = ((\xi_{ij}))_{1 \leq i, j \leq n}$, ξ_{ij} i.i.d $N_{\mathbb{C}}(0, 1)$ (NO normalization by \sqrt{n})
- $n = \infty$: $\mu = \lim_{n \rightarrow \infty} \mu_n$ (Ginibre ensemble)

Ginibre Ensemble

- Finite n : $\mu_n =$ Eigenvalues of $G_n = ((\xi_{ij}))_{1 \leq i, j \leq n}$, ξ_{ij} i.i.d $N_{\mathbb{C}}(0, 1)$ (NO normalization by \sqrt{n})
- $n = \infty$: $\mu = \lim_{n \rightarrow \infty} \mu_n$ (Ginibre ensemble)
- Translation Invariant (in fact Ergodic)

Gaussian Analytic Function

- Finite n : $f_n(z) = \xi_0 + \frac{\xi_1}{\sqrt{1!}}z + \dots + \frac{\xi_k}{\sqrt{k!}}z^k + \dots + \frac{\xi_n}{\sqrt{n!}}z^n$
 $\nu_n = \text{Zeroes of } f_n \quad (\xi_i \text{ iid } N_{\mathbb{C}}(0, 1))$

Gaussian Analytic Function

- Finite n : $f_n(z) = \xi_0 + \frac{\xi_1}{\sqrt{1!}}z + \dots + \frac{\xi_k}{\sqrt{k!}}z^k + \dots + \frac{\xi_n}{\sqrt{n!}}z^n$
 $\nu_n = \text{Zeroes of } f_n \quad (\xi_i \text{ iid } N_{\mathbb{C}}(0, 1))$
- $n = \infty$:
 - $\nu = \lim_{n \rightarrow \infty} \nu_n$
 - Zeroes of $f(z) = \sum_{k=0}^{\infty} \frac{\xi_k}{\sqrt{k!}}z^k$

Gaussian Analytic Function

- Finite n : $f_n(z) = \xi_0 + \frac{\xi_1}{\sqrt{1!}}z + \dots + \frac{\xi_k}{\sqrt{k!}}z^k + \dots + \frac{\xi_n}{\sqrt{n!}}z^n$
 $\nu_n = \text{Zeroes of } f_n \quad (\xi_i \text{ iid } N_{\mathbb{C}}(0, 1))$
- $n = \infty$:
 - $\nu = \lim_{n \rightarrow \infty} \nu_n$
 - Zeroes of $f(z) = \sum_{k=0}^{\infty} \frac{\xi_k}{\sqrt{k!}}z^k$
- ν is **Translation Invariant** (and Ergodic)

Gaussian Analytic Function

- Finite n : $f_n(z) = \xi_0 + \frac{\xi_1}{\sqrt{1!}}z + \dots + \frac{\xi_k}{\sqrt{k!}}z^k + \dots + \frac{\xi_n}{\sqrt{n!}}z^n$
 $\nu_n = \text{Zeroes of } f_n \quad (\xi_i \text{ iid } N_{\mathbb{C}}(0, 1))$
- $n = \infty$:
 - $\nu = \lim_{n \rightarrow \infty} \nu_n$
 - Zeroes of $f(z) = \sum_{k=0}^{\infty} \frac{\xi_k}{\sqrt{k!}}z^k$
- ν is **Translation Invariant** (and Ergodic)

Theorem (Sodin rigidity)

$f(z)$ is the unique (up to a deterministic multiplier) Gaussian entire function with a translation invariant zero process of intensity 1.

- We consider the conditional distribution $\rho_\omega(\zeta)$ of the points (denoted by ζ) inside a disk \mathbb{D} given the points outside \mathbb{D} (denoted by ω)

- We consider the conditional distribution $\rho_\omega(\zeta)$ of the points (denoted by ζ) inside a disk \mathbb{D} given the points outside \mathbb{D} (denoted by ω)
- In Poisson point process, the points inside and outside \mathbb{D} are independent of each other

Our Results: Ginibre Ensemble

Theorem (G.,Peres)

In the Ginibre ensemble,

(i) The points ω outside \mathbb{D} determine exactly the Number $N(\omega)$ of the points ζ inside \mathbb{D}

Our Results: Ginibre Ensemble

Theorem (G.,Peres)

In the Ginibre ensemble,

*(i) The points ω outside \mathbb{D} **determine** exactly the **Number $N(\omega)$** of the points ζ inside \mathbb{D} “and nothing more”*

Theorem (G.,Peres)

In the Ginibre ensemble,

(i) The points ω outside \mathbb{D} determine exactly the Number $N(\omega)$ of the points ζ inside \mathbb{D} “and nothing more”

(ii) A.e. ω ,

(a) The conditional measure $\rho_\omega(\zeta)$ is absolutely continuous wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$,

Theorem (G.,Peres)

In the Ginibre ensemble,

*(i) The points ω outside \mathbb{D} **determine** exactly the **Number $N(\omega)$** of the points ζ inside \mathbb{D} “and nothing more”*

(ii) A.e. ω ,

*(a) The conditional measure $\rho_\omega(\zeta)$ is **absolutely continuous wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$** , and hence has a probability density function $f_\omega(\zeta)$*

Theorem (G.,Peres)

In the Ginibre ensemble,

(i) The points ω outside \mathbb{D} determine exactly the Number $N(\omega)$ of the points ζ inside \mathbb{D} “and nothing more”

(ii) A.e. ω ,

(a) The conditional measure $\rho_\omega(\zeta)$ is absolutely continuous wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$, and hence has a probability density function $f_\omega(\zeta)$

(b) $f_\omega(\zeta) > 0$ a.e. wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$

Theorem (G.,Peres)

In the Ginibre ensemble,

(i) *The points ω outside \mathbb{D} determine exactly the Number $N(\omega)$ of the points ζ inside \mathbb{D} “and nothing more”*

(ii) *A.e. ω ,*

(a) *The conditional measure $\rho_\omega(\zeta)$ is absolutely continuous wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$, and hence has a probability density function $f_\omega(\zeta)$*

(b) *$f_\omega(\zeta) > 0$ a.e. wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$*

(c) *$m(\omega)|\Delta(\zeta)|^2 \leq f_\omega(\zeta) \leq M(\omega)|\Delta(\zeta)|^2$*

Theorem (G.,Peres)

In the Ginibre ensemble,

(i) The points ω outside \mathbb{D} *determine* exactly the *Number* $N(\omega)$ of the points ζ inside \mathbb{D} “and nothing more”

(ii) A.e. ω ,

(a) The conditional measure $\rho_\omega(\zeta)$ is *absolutely continuous wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$* , and hence has a probability density function $f_\omega(\zeta)$

(b) $f_\omega(\zeta) > 0$ a.e. wrt Lebesgue measure on $\mathbb{D}^{N(\omega)}$

(c) $m(\omega)|\Delta(\zeta)|^2 \leq f_\omega(\zeta) \leq M(\omega)|\Delta(\zeta)|^2$

$M(\omega)$ and $m(\omega)$ positive constants

$\Delta(\zeta) = \prod_{i < j} (\zeta_i - \zeta_j)$ (Vandermonde)

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .
- E.g. :
 - Ginibre ensemble

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .
- E.g. :
 - Ginibre ensemble
 - Finite point process (fixed size n) - e.g. eigenvalues of a $n \times n$ random matrix

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .
- E.g. :
 - Ginibre ensemble
 - Finite point process (fixed size n) - e.g. eigenvalues of a $n \times n$ random matrix
- Non e.g.: Poisson point process is NOT rigid

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .
- E.g. :
 - Ginibre ensemble
 - Finite point process (fixed size n) - e.g. eigenvalues of a $n \times n$ random matrix
- Non e.g.: Poisson point process is NOT rigid
- Rigidity does Not pass to the Limit!

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .
- E.g. :
 - Ginibre ensemble
 - Finite point process (fixed size n) - e.g. eigenvalues of a $n \times n$ random matrix
- Non e.g.: Poisson point process is NOT rigid
- Rigidity does Not pass to the Limit!

Take n points in a disk of area n . Finite, rigid.

- **Rigidity** A process is said to be rigid if, given the points outside a domain D , we can predict the exact number of points in \mathbb{D} with probability 1 .
- E.g. :
 - Ginibre ensemble
 - Finite point process (fixed size n) - e.g. eigenvalues of a $n \times n$ random matrix
- Non e.g.: Poisson point process is NOT rigid
- Rigidity does Not pass to the Limit!

Take n points in a disk of area n . Finite, rigid.
Limit as $n \rightarrow \infty$ is Poisson : Not Rigid !

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

*(i) The points ω outside \mathbb{D} **determine** exactly the **Number $N(\omega)$***

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

*(i) The points ω outside \mathbb{D} **determine** exactly the **Number** $N(\omega)$ and **Sum** $S(\omega)$ of points ζ inside \mathbb{D} ,*

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

*(i) The points ω outside \mathbb{D} **determine** exactly the **Number** $N(\omega)$ and **Sum** $S(\omega)$ of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$),*

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

*(i) The points ω outside \mathbb{D} **determine** exactly the **Number $N(\omega)$** and **Sum $S(\omega)$** of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$), “and nothing more”*

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

(i) The points ω outside \mathbb{D} *determine* exactly the *Number* $N(\omega)$ and *Sum* $S(\omega)$ of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$), “and nothing more”

(ii) A.e. ω ,

(a) $\rho_\omega(\zeta)$ is *absolutely continuous wrt Lebesgue measure* on $\Sigma_{S(\omega)}$,

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

(i) The points ω outside \mathbb{D} *determine* exactly the *Number* $N(\omega)$ and *Sum* $S(\omega)$ of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$), “and nothing more”

(ii) A.e. ω ,

(a) $\rho_\omega(\zeta)$ is *absolutely continuous wrt Lebesgue measure* on $\Sigma_{S(\omega)}$, and hence has a probability density function $f_\omega(\zeta)$

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

(i) The points ω outside \mathbb{D} *determine* exactly the *Number* $N(\omega)$ and *Sum* $S(\omega)$ of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$), “and nothing more”

(ii) A.e. ω ,

(a) $\rho_\omega(\zeta)$ is *absolutely continuous wrt Lebesgue measure* on $\Sigma_{S(\omega)}$, and hence has a probability density function $f_\omega(\zeta)$

(b) $f_\omega(\zeta) > 0$ a.e. wrt Lebesgue measure on $\Sigma_{S(\omega)}$

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

(i) The points ω outside \mathbb{D} *determine* exactly the *Number* $N(\omega)$ and *Sum* $S(\omega)$ of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$), “and nothing more”

(ii) A.e. ω ,

(a) $\rho_\omega(\zeta)$ is *absolutely continuous wrt Lebesgue measure on* $\Sigma_{S(\omega)}$, and hence has a probability density function $f_\omega(\zeta)$

(b) $f_\omega(\zeta) > 0$ a.e. wrt Lebesgue measure on $\Sigma_{S(\omega)}$

(c) $m(\omega)|\Delta(\zeta)|^2 \leq f_\omega(\zeta) \leq M(\omega)|\Delta(\zeta)|^2$ a.e.

Our Results: Zeroes of Gaussian Analytic Function

Theorem (G., Peres)

In the GAF Zero ensemble,

(i) The points ω outside \mathbb{D} *determine* exactly the *Number* $N(\omega)$ and *Sum* $S(\omega)$ of points ζ inside \mathbb{D} , (so $\zeta \in$ constant sum hypersurface $\Sigma_{S(\omega)} \subset \mathbb{D}^{N(\omega)}$), “and nothing more”

(ii) A.e. ω ,

(a) $\rho_\omega(\zeta)$ is *absolutely continuous wrt Lebesgue measure on* $\Sigma_{S(\omega)}$, and hence has a probability density function $f_\omega(\zeta)$

(b) $f_\omega(\zeta) > 0$ a.e. wrt Lebesgue measure on $\Sigma_{S(\omega)}$

(c) $m(\omega)|\Delta(\zeta)|^2 \leq f_\omega(\zeta) \leq M(\omega)|\Delta(\zeta)|^2$ a.e.

$\Sigma_{S(\omega)}$: constant sum hypersurface $\sum_{i=1}^{N(\omega)} \zeta_i = S(\omega)$ inside $\mathbb{D}^{N(\omega)}$

$M(\omega)$, $m(\omega)$ and $\Delta(\zeta)$ are as before.

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$
- **Scaling:** $\varphi_L(z) = \varphi\left(\frac{z}{L}\right)$

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$
- **Scaling:** $\varphi_L(z) = \varphi\left(\frac{z}{L}\right)$
- **(Sodin Tsirelson)** $\text{var}[\int \varphi_L d\nu] = O\left(\frac{1}{L^2}\right)$

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$
- **Scaling:** $\varphi_L(z) = \varphi\left(\frac{z}{L}\right)$
- **(Sodin Tsirelson)** $\text{var}[\int \varphi_L d\nu] = O\left(\frac{1}{L^2}\right)$
- Take φ as roughly $1_{\mathbb{D}}$ (e.g. $1_{\mathbb{D}} \leq \varphi \leq 1_{2\mathbb{D}}, \varphi \in C_c^2$)

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$
- **Scaling:** $\varphi_L(z) = \varphi\left(\frac{z}{L}\right)$
- **(Sodin Tsirelson)** $\text{var}[\int \varphi_L d\nu] = O\left(\frac{1}{L^2}\right)$
- Take φ as roughly $1_{\mathbb{D}}$ (e.g. $1_{\mathbb{D}} \leq \varphi \leq 1_{2\mathbb{D}}, \varphi \in C_c^2$)
- By Sodin Tsirelson,
 $\int \varphi_L d\nu \approx \mathbb{E}[\int \varphi_L d\nu] = \int \varphi_L(z) \rho_1(z) dm(z)$

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$
- **Scaling:** $\varphi_L(z) = \varphi\left(\frac{z}{L}\right)$
- **(Sodin Tsirelson)** $\text{var}[\int \varphi_L d\nu] = O\left(\frac{1}{L^2}\right)$
- Take φ as roughly $1_{\mathbb{D}}$ (e.g. $1_{\mathbb{D}} \leq \varphi \leq 1_{2\mathbb{D}}, \varphi \in C_c^2$)
- By Sodin Tsirelson,
 $\int \varphi_L d\nu \approx \mathbb{E}[\int \varphi_L d\nu] = \int \varphi_L(z) \rho_1(z) dm(z)$
- But $\int \varphi_L d\nu = n(\mathbb{D}) + \int_{\mathbb{D}_L \setminus \mathbb{D}} \varphi_L d\nu$

Proof Sketch: Rigidity of Number of Points of GAF

- Given: outside zeroes of GAF
Want: number of inside zeroes
- **Linear Statistic:** $\int \varphi d\nu, \varphi \in C_c^2(\mathbb{C})$
- **Scaling:** $\varphi_L(z) = \varphi\left(\frac{z}{L}\right)$
- **(Sodin Tsirelson)** $\text{var}[\int \varphi_L d\nu] = O\left(\frac{1}{L^2}\right)$
- Take φ as roughly $1_{\mathbb{D}}$ (e.g. $1_{\mathbb{D}} \leq \varphi \leq 1_{2\mathbb{D}}, \varphi \in C_c^2$)
- By Sodin Tsirelson,
$$\int \varphi_L d\nu \approx \mathbb{E}[\int \varphi_L d\nu] = \int \varphi_L(z) \rho_1(z) dm(z)$$
- But $\int \varphi_L d\nu = n(\mathbb{D}) + \int_{\mathbb{D}_L \setminus \mathbb{D}} \varphi_L d\nu$
- Know outside zeroes \Rightarrow Know $\int_{\mathbb{D}_L \setminus \mathbb{D}} \varphi_L d\nu \Rightarrow$ Compute $n(\mathbb{D})$ approximately, now let $L \rightarrow \infty$

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.
- For Poisson point process, it is known that:
 - $\exists 0 < r_{\text{crit}} < \infty$ such that for $r < r_{\text{crit}}$, there is no infinite cluster a.s., and for $r > r_{\text{crit}}$, there is an infinite cluster a.s.

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.
- For Poisson point process, it is known that:
 - $\exists 0 < r_{\text{crit}} < \infty$ such that for $r < r_{\text{crit}}$, there is no infinite cluster a.s., and for $r > r_{\text{crit}}$, there is an infinite cluster a.s.
 - The infinite cluster, when it exists, is unique.

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.
- For Poisson point process, it is known that:
 - $\exists 0 < r_{\text{crit}} < \infty$ such that for $r < r_{\text{crit}}$, there is no infinite cluster a.s., and for $r > r_{\text{crit}}$, there is an infinite cluster a.s.
 - The infinite cluster, when it exists, is unique.
- Proofs use spatial independence in a strong way : what about correlated point processes?

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.
- For Poisson point process, it is known that:
 - $\exists 0 < r_{\text{crit}} < \infty$ such that for $r < r_{\text{crit}}$, there is no infinite cluster a.s., and for $r > r_{\text{crit}}$, there is an infinite cluster a.s.
 - The infinite cluster, when it exists, is unique.
- Proofs use spatial independence in a strong way : what about correlated point processes? No “finite energy” either !

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.
- For Poisson point process, it is known that:
 - $\exists 0 < r_{\text{crit}} < \infty$ such that for $r < r_{\text{crit}}$, there is no infinite cluster a.s., and for $r > r_{\text{crit}}$, there is an infinite cluster a.s.
 - The infinite cluster, when it exists, is unique.
- Proofs use spatial independence in a strong way : what about correlated point processes? No “finite energy” either !

Theorem (G., Krishnapur, Peres)

For Gaussian zeroes and Ginibre eigenvalues, we establish both phase transition and uniqueness.

Continuum percolation

- Put a disk of deterministic radius r around each point of a point process π .
- Two points are neighbours \iff their disks intersect.
- For Poisson point process, it is known that:
 - $\exists 0 < r_{\text{crit}} < \infty$ such that for $r < r_{\text{crit}}$, there is no infinite cluster a.s., and for $r > r_{\text{crit}}$, there is an infinite cluster a.s.
 - The infinite cluster, when it exists, is unique.
- Proofs use spatial independence in a strong way : what about correlated point processes? No “finite energy” either !

Theorem (G., Krishnapur, Peres)

For Gaussian zeroes and Ginibre eigenvalues, we establish both phase transition and uniqueness.

Uniqueness involves understanding of rigidity and tolerance.

Theorem (Reconstruction of Gaussian Analytic Function, G.-Peres)

The zeroes of the GAF determine the function a.s. (up to a multiplicative factor of modulus 1). In other words, if ν denotes the zeroes of the GAF f , then \exists an analytic function

$$g(z) = \sum_{k=0}^{\infty} a_k(\nu) z^k \text{ such that } f(z) = \gamma \cdot g(z)$$

Here γ follows $\text{Unif}(S^1)$ and is independent of ν .

Some Interesting Results en route

Theorem (Reconstruction of Gaussian Analytic Function, G.-Peres)

The zeroes of the GAF determine the function a.s. (up to a multiplicative factor of modulus 1). In other words, if ν denotes the zeroes of the GAF f , then \exists an analytic function

$$g(z) = \sum_{k=0}^{\infty} a_k(\nu) z^k \text{ such that } f(z) = \gamma \cdot g(z)$$

Here γ follows $\text{Unif}(S^1)$ and is independent of ν .

Theorem (Stability of Inverse Sums, G.-Peres)

Let z denote the points of Ginibre ensemble. Then the random series $\sum_{|z| \uparrow} \frac{1}{z}$ converges almost surely, and in fact, has finite first moment.

- Completeness of random exponential systems

- Completeness of random exponential systems
- A conjecture of Lyons-Steif

- Completeness of random exponential systems
- A conjecture of Lyons-Steif
- Studying mixtures of point processes

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$
- $\text{Span}(\mathcal{E}_\Lambda) \subset L_2(-\pi, \pi)$.

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$
- $\text{Span}(\mathcal{E}_\Lambda) \subset L_2(-\pi, \pi)$.
- Question: Is $\text{Span}(\mathcal{E}_\Lambda) = L_2(-\pi, \pi)$?

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$
- $\text{Span}(\mathcal{E}_\Lambda) \subset L_2(-\pi, \pi)$.
- Question: Is $\text{Span}(\mathcal{E}_\Lambda) = L_2(-\pi, \pi)$?
- Classical question in harmonic analysis (Levinson, Redheffer, Beurling, Malliavin,)

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$
- $\text{Span}(\mathcal{E}_\Lambda) \subset L_2(-\pi, \pi)$.
- Question: Is $\text{Span}(\mathcal{E}_\Lambda) = L_2(-\pi, \pi)$?
- Classical question in harmonic analysis (Levinson, Redheffer, Beurling, Malliavin,)
- Classical results are parametrized by some sort of asymptotic density of Λ ; complete if supercritical, incomplete if subcritical.

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$
- $\text{Span}(\mathcal{E}_\Lambda) \subset L_2(-\pi, \pi)$.
- Question: Is $\text{Span}(\mathcal{E}_\Lambda) = L_2(-\pi, \pi)$?
- Classical question in harmonic analysis (Levinson, Redheffer, Beurling, Malliavin,)
- Classical results are parametrized by some sort of asymptotic density of Λ ; complete if supercritical, incomplete if subcritical.
- Tricky situation at criticality (density 1).

Completeness of random exponential systems

- For a discrete point set $\Lambda \subset \mathbb{R}$, consider the space of exponential functions $\mathcal{E}_\Lambda := \{e_\lambda(x) = e^{i\lambda x} : \lambda \in \Lambda\}$
- $\text{Span}(\mathcal{E}_\Lambda) \subset L_2(-\pi, \pi)$.
- Question: Is $\text{Span}(\mathcal{E}_\Lambda) = L_2(-\pi, \pi)$?
- Classical question in harmonic analysis (Levinson, Redheffer, Beurling, Malliavin,)
- Classical results are parametrized by some sort of asymptotic density of Λ ; complete if supercritical, incomplete if subcritical.
- Tricky situation at criticality (density 1).
- What if pathological configurations are eliminated by choosing a “generic” point configuration - i.e., a point process ?

Theorem (G.)

Let Π be a determinantal point process with kernel K and background measure μ , such that K is the integral kernel corresponding to a projection on to a subspace \mathcal{H} of $L_2(\mu)$. Clearly, $\mathcal{E}_\Pi = \{K(x, \cdot) : x \in \Pi\} \subset \mathcal{H}$.

Theorem (G.)

Let Π be a determinantal point process with kernel K and background measure μ , such that K is the integral kernel corresponding to a projection on to a subspace \mathcal{H} of $L_2(\mu)$.

Clearly, $\mathcal{E}_\Pi = \{K(x, \cdot) : x \in \Pi\} \subset \mathcal{H}$.

If Π is rigid, then we have $\text{Span}(\mathcal{E}_\Pi) = \mathcal{H}$ with probability 1.

Theorem (G.)

Let Π be a determinantal point process with kernel K and background measure μ , such that K is the integral kernel corresponding to a projection on to a subspace \mathcal{H} of $L_2(\mu)$.

Clearly, $\mathcal{E}_\Pi = \{K(x, \cdot) : x \in \Pi\} \subset \mathcal{H}$.

If Π is rigid, then we have $\text{Span}(\mathcal{E}_\Pi) = \mathcal{H}$ with probability 1.

- Theorem + Fourier transform gives positive answer to the completeness question for Λ sampled from the sine kernel process.

Theorem (G.)

Let Π be a determinantal point process with kernel K and background measure μ , such that K is the integral kernel corresponding to a projection on to a subspace \mathcal{H} of $L_2(\mu)$.

Clearly, $\mathcal{E}_\Pi = \{K(x, \cdot) : x \in \Pi\} \subset \mathcal{H}$.

If Π is rigid, then we have $\text{Span}(\mathcal{E}_\Pi) = \mathcal{H}$ with probability 1.

- Theorem + Fourier transform gives positive answer to the completeness question for Λ sampled from the sine kernel process.
- Theorem implies positive answer to analogous completeness question in 2-d for random exponentials (sampled from the Ginibre ensemble) inside the Fock Bargmann space.

A Hierarchy of Point Processes

- π is **rigid** at level k if
 - The points of π outside \mathbb{D} determine $0, 1, \dots, (k - 1)$ moments of the points in \mathbb{D}

A Hierarchy of Point Processes

- π is **rigid** at level k if
 - The points of π outside \mathbb{D} determine $0, 1, \dots, (k - 1)$ moments of the points in \mathbb{D}
 - Conditional distribution of the points of π inside \mathbb{D} given the outside has a non-vanishing density wrt Lebesgue measure on the submanifold of $\mathbb{D}^{N(\omega)}$ defined by $0, 1, \dots, (k - 1)$ moments being conserved.

A Hierarchy of Point Processes

- π is **rigid** at level k if
 - The points of π outside \mathbb{D} determine $0, 1, \dots, (k - 1)$ moments of the points in \mathbb{D}
 - Conditional distribution of the points of π inside \mathbb{D} given the outside has a non-vanishing density wrt Lebesgue measure on the submanifold of $\mathbb{D}^{N(\omega)}$ defined by $0, 1, \dots, (k - 1)$ moments being conserved.
- Poisson is at **Level 0** (i.e. **no moments** conserved)

A Hierarchy of Point Processes

- π is **rigid** at level k if
 - The points of π outside \mathbb{D} determine $0, 1, \dots, (k - 1)$ moments of the points in \mathbb{D}
 - Conditional distribution of the points of π inside \mathbb{D} given the outside has a non-vanishing density wrt Lebesgue measure on the submanifold of $\mathbb{D}^{N(\omega)}$ defined by $0, 1, \dots, (k - 1)$ moments being conserved.
- Poisson is at **Level 0** (i.e. **no moments** conserved)
- Ginibre is at **Level 1** (i.e. **only 0th moment** conserved)

A Hierarchy of Point Processes

- π is **rigid** at level k if
 - The points of π outside \mathbb{D} determine $0, 1, \dots, (k - 1)$ moments of the points in \mathbb{D}
 - Conditional distribution of the points of π inside \mathbb{D} given the outside has a non-vanishing density wrt Lebesgue measure on the submanifold of $\mathbb{D}^{N(\omega)}$ defined by $0, 1, \dots, (k - 1)$ moments being conserved.
- Poisson is at **Level 0** (i.e. **no moments** conserved)
- Ginibre is at **Level 1** (i.e. **only 0th moment** conserved)
- GAF Zeros is at **Level 2** (i.e. **only 0th and 1st moment** conserved)

A Hierarchy of Point Processes

- π is **rigid** at level k if
 - The points of π outside \mathbb{D} determine $0, 1, \dots, (k - 1)$ moments of the points in \mathbb{D}
 - Conditional distribution of the points of π inside \mathbb{D} given the outside has a non-vanishing density wrt Lebesgue measure on the submanifold of $\mathbb{D}^{N(\omega)}$ defined by $0, 1, \dots, (k - 1)$ moments being conserved.
- Poisson is at **Level 0** (i.e. **no moments** conserved)
- Ginibre is at **Level 1** (i.e. **only 0th moment** conserved)
- GAF Zeros is at **Level 2** (i.e. **only 0th and 1st moment** conserved)
- Natural point processes for Levels $k \geq 3$??

- **k rigidity** There is a family of analytic functions with Gaussian coefficients such that the points outside a disk determine the first k moments of the points inside, and “nothing more”.

- **k rigidity** There is a family of analytic functions with Gaussian coefficients such that the points outside a disk determine the first k moments of the points inside, and “nothing more”.
- **Rigidity for dpp** If a determinantal point process has a kernel which is a proper contraction, then it must be insertion and deletion tolerant.

Thank you !!